

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

The following is a list of recommended preventive services and corresponding CPT, HCPCS, and diagnoses codes to assist Medicaid health plans and providers in identifying Medicaid covered preventive services. Providers are encouraged to identify the service as preventive by including HCPCS modifier 33, preventive service. In addition to modifier 33, a diagnosis may be required to support that the service was preventive.

This information is provided as a reference only and not intended to replace the clinical judgement of the providers. Other coverage determinations may apply.

Additional information about preventive care guidelines is available at the following:

- United States Preventive Services Task Force grade A and B services
<http://www.uspreventiveservicestaskforce.org/Page/Name/recommendations>
- Advisory Committee on Immunization Practices recommended vaccines
<http://www.cdc.gov/vaccines/hcp/acip-recs/index.html>
- Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine recommended preventive services for women
<http://www.nationalacademies.org/hmd/Reports/2011/Clinical-Preventive-Services-for-Women-Closing-the-Gaps.aspx>
- Early and Periodic Screening, Diagnosis and Treatment services as defined in the current periodicity schedule by the American Academy of Pediatrics
http://brightfutures.aap.org/clinical_practice.html

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Abdominal aortic aneurysm screening: men	The USPSTF recommends one-time screening for abdominal aortic aneurysm by ultrasonography in men ages 65 to 75 years who have ever smoked.	76706, 76770*, 76775*	Z136, Z87891
Alcohol misuse: screening and counseling	The USPSTF recommends that clinicians screen adults age 18 years or older for alcohol misuse and provide persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce alcohol misuse.	99408, 99409	Reason For Visit

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Aspirin preventive medication: adults aged 50 to 59 years with a $\geq 10\%$ 10-year cardiovascular risk	The USPSTF recommends initiating low-dose aspirin use for the primary prevention of cardiovascular disease and colorectal cancer in adults aged 50 to 59 years who have a 10% or greater 10-year cardiovascular risk, are not at increased risk for bleeding, have a life expectancy of at least 10 years, and are willing to take low-dose aspirin daily for at least 10 years.	Pharmacy Benefit	NA
Bacteriuria screening: pregnant women	The USPSTF recommends screening for asymptomatic bacteriuria with urine culture in pregnant women at 12 to 16 weeks' gestation or at the first prenatal visit, if later.	87086*, 87088*	Reason For Screen
Blood pressure screening: adults	The USPSTF recommends screening for high blood pressure in adults aged 18 years or older. The USPSTF recommends obtaining measurements outside of the clinical setting for diagnostic confirmation before starting treatment.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit
BRCA risk assessment and genetic counseling/testing	The USPSTF recommends that primary care providers screen women who have family members with breast, ovarian, tubal, or peritoneal cancer with one of several screening tools designed to identify a family history that may be associated with an increased risk for potentially harmful mutations in breast cancer susceptibility genes (BRCA1 or BRCA2). Women with positive screening results should receive genetic counseling and, if indicated after counseling, BRCA testing.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Breast cancer preventive medications	The USPSTF recommends that clinicians engage in shared, informed decision-making with women who are at increased risk for breast cancer about medications to reduce their risk. For women who are at increased risk for breast cancer and at low risk for adverse medication effects, clinicians should offer to prescribe risk-reducing medications, such as tamoxifen or raloxifene.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Breast cancer screening	The USPSTF recommends screening mammography for women, with or without clinical breast examination, every 1 to 2 years for women age 40 years and older.	77063, 77067	Reason For Screen
Breastfeeding interventions	The USPSTF recommends providing interventions during pregnancy and after birth to support breastfeeding.	S9443	Reason For Visit
Cervical cancer screening	The USPSTF recommends screening for cervical cancer in women ages 21 to 65 years with cytology (Pap smear) every 3 years or, for women ages 30 to 65 years who want to lengthen the screening interval, screening with a combination of cytology and human papillomavirus (HPV) testing every 5 years.	87623-87625, 88141-88148, 88155, 88164-88167, 88174-88175, G0101, G0476, Q0091	Reason For Screen
Chlamydia screening: women	The USPSTF recommends screening for chlamydia in sexually active women age 24 years or younger and in older women who are at increased risk for infection.	87110, 87270, 87320, 87490-87492, 87810	Reason For Screen
Colorectal cancer screening	The USPSTF recommends screening for colorectal cancer starting at age 50 years and continuing until age 75 years.	00812, 45330*, 45331*, 45333*, 45338*, 45346*, 45378*, 45380, 45384*, 45385*, 45388*, 81528, 82270, 82274*, G0104, G0105, G0121, G0328,	Z1211, Z1212, Z1213, Z800, Z8371, Z86010
Dental caries prevention: infants and children up to age 5 years	The USPSTF recommends the application of fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption in primary care practices. The USPSTF recommends primary care clinicians prescribe oral fluoride supplementation starting at age 6 months for children whose water supply is fluoride deficient.	99188, D1206, D1208	Reason For Visit
Depression screening: adolescents	The USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12 to 18 years. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.	96127	Reason For Visit

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Depression screening: adults	The USPSTF recommends screening for depression in the general adult population, including pregnant and postpartum women. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.	96127*	Reason For Visit
Diabetes screening	The USPSTF recommends screening for abnormal blood glucose as part of cardiovascular risk assessment in adults aged 40 to 70 years who are overweight or obese. Clinicians should offer or refer patients with abnormal blood glucose to intensive behavioral counseling interventions to promote a healthful diet and physical activity.	82947*, 82948*, 82950*, 82951*, 82952*, 83036*	Z131, Z833
Falls prevention in older adults: exercise or physical therapy	The USPSTF recommends exercise or physical therapy to prevent falls in community-dwelling adults age 65 years and older who are at increased risk for falls.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit
Falls prevention in older adults: vitamin D	The USPSTF recommends vitamin D supplementation to prevent falls in community-dwelling adults age 65 years and older who are at increased risk for falls.	Pharmacy Benefit	NA
Folic acid supplementation	The USPSTF recommends that all women who are planning or capable of pregnancy take a daily supplement containing 0.4 to 0.8 mg (400 to 800 µg) of folic acid.	Pharmacy Benefit	NA
Gestational diabetes mellitus screening	The USPSTF recommends screening for gestational diabetes mellitus in asymptomatic pregnant women after 24 weeks of gestation.	82950*, 82951*, 82952*	Reason For Screen
Gonorrhea prophylactic medication: newborns	The USPSTF recommends prophylactic ocular topical medication for all newborns for the prevention of gonococcal ophthalmia neonatorum.	Newborn Inpatient	Reason For Visit
Gonorrhea screening: women	The USPSTF recommends screening for gonorrhea in sexually active women age 24 years or younger and in older women who are at increased risk for infection.	87590, 87591, 87592, 87850,	Reason For Screen

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Healthy diet and physical activity counseling to prevent cardiovascular disease: adults with cardiovascular risk factors	The USPSTF recommends offering or referring adults who are overweight or obese and have additional cardiovascular disease (CVD) risk factors to intensive behavioral counseling interventions to promote a healthful diet and physical activity for CVD prevention.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit
Hearing loss screening: newborns	The USPSTF recommends screening for hearing loss in all newborn infants.	Inpatient Newborn Screen 92551-92553, 92558, 92568, 92585-92588, V5008	Reason For Visit
Hemoglobinopathies screening: newborns	The USPSTF recommends screening for sickle cell disease in newborns.	Inpatient Newborn Screen	Reason For Visit
Hepatitis B screening: nonpregnant adolescents and adults	The USPSTF recommends screening for hepatitis B virus infection in persons at high risk for infection.	86704-86706, 87340, 87341, G0499	Reason For Screen
Hepatitis B screening: pregnant women	The USPSTF strongly recommends screening for hepatitis B virus infection in pregnant women at their first prenatal visit.	80055,80081, 86704-86706, 87340, 87341, G0499	Reason For Screen
Hepatitis C virus infection screening: adults	The USPSTF recommends screening for hepatitis C virus (HCV) infection in persons at high risk for infection. The USPSTF also recommends offering one-time screening for HCV infection to adults born between 1945 and 1965.	86803, G0472	Reason For Screen
HIV screening: nonpregnant adolescents and adults	The USPSTF recommends that clinicians screen for HIV infection in adolescents and adults ages 15 to 65 years. Younger adolescents and older adults who are at increased risk should also be screened.	86689*, 86701*, 86702*, 86703*, 87389*, 87390*, 87391*, 87534*, 87535*, 87536*, 87537*, 87538*, 87539*, 87806*, G0432, G0433, G0435	Z114

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
HIV screening: pregnant women	The USPSTF recommends that clinicians screen all pregnant women for HIV, including those who present in labor who are untested and whose HIV status is unknown.	80081*, 86689*, 86701*, 86702*, 86703*, 87389*, 87390*, 87391*, 87534*, 87535*, 87536*, 87537*, 87538*, 87539*, 87806*, G0432, G0433, G0435	Reason For Screen
Hypothyroidism screening: newborns	The USPSTF recommends screening for congenital hypothyroidism in newborns.	Inpatient Newborn Screen	Reason For Visit
Intimate partner violence screening: women of childbearing age	The USPSTF recommends that clinicians screen women of childbearing age for intimate partner violence, such as domestic violence, and provide or refer women who screen positive to intervention services. This recommendation applies to women who do not have signs or symptoms of abuse.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Lung cancer screening	The USPSTF recommends annual screening for lung cancer with low-dose computed tomography in adults ages 55 to 80 years who have a 30 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.	71250*, G0297	F17200-F17299, Z122, Z720, Z87891,
Obesity screening and counseling: adults	The USPSTF recommends screening all adults for obesity. Clinicians should offer or refer patients with a body mass index of 30 kg/m ² or higher to intensive, multicomponent behavioral interventions.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Obesity screening: children and adolescents	The USPSTF recommends that clinicians screen for obesity in children and adolescents 6 years and older and offer or refer them to comprehensive, intensive behavioral interventions to promote improvements in weight status.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Osteoporosis screening: women	The USPSTF recommends screening for osteoporosis in women age 65 years and older and in younger women whose fracture risk is equal to or greater than that of a 65-year-old white woman who has no additional risk factors.	76977*, 77078*, 77080*, 77081*	Z13820, Z8262
Phenylketonuria screening: newborns	The USPSTF recommends screening for phenylketonuria in newborns.	Inpatient Newborn Screen	Reason For Visit
Preeclampsia prevention: aspirin	The USPSTF recommends the use of low-dose aspirin (81 mg/d) as preventive medication after 12 weeks of gestation in women who are at high risk for preeclampsia.	Pharmacy Benefit	NA
Preeclampsia: screening	The USPSTF recommends screening for preeclampsia in pregnant women with blood pressure measurements throughout pregnancy.	Covered during preventive medicine services or a focused E/M visit	Reason For Visit
Rh incompatibility screening: 24–28 weeks' gestation	The USPSTF recommends repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24 to 28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.	80055*, 80081*, 86901*	Reason For Screen
Rh incompatibility screening: first pregnancy visit	The USPSTF strongly recommends Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.	80055*, 80081*, 86901*	Reason For Screen
Sexually transmitted infections counseling	The USPSTF recommends intensive behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit
Skin cancer behavioral counseling	The USPSTF recommends counseling children, adolescents, and young adults ages 10 to 24 years who have fair skin about minimizing their exposure to ultraviolet radiation to reduce risk for skin cancer.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Statin preventive medication: adults ages 40–75 years with no history of CVD, 1 or more CVD risk factors, and a calculated 10-year CVD event risk of 10% or greater	The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (i.e., symptomatic coronary artery disease or ischemic stroke) use a low- to moderate-dose statin for the prevention of CVD events and mortality when all of the following criteria are met: 1) they are ages 40 to 75 years; 2) they have 1 or more CVD risk factors (i.e., dyslipidemia, diabetes, hypertension, or smoking); and 3) they have a calculated 10-year risk of a cardiovascular event of 10% or greater. Identification of dyslipidemia and calculation of 10-year CVD event risk requires universal lipids screening in adults ages 40 to 75 years.	Pharmacy Benefit	NA
Syphilis screening: nonpregnant persons	The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection.	86592-86593	Reason For Screen
Syphilis screening: pregnant women	The USPSTF recommends that clinicians screen all pregnant women for syphilis infection.	80055, 80081, 86592, 86593	Reason For Screen
Tobacco use counseling and interventions: nonpregnant adults	The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and U.S. Food and Drug Administration (FDA)–approved pharmacotherapy for cessation to adults who use tobacco.	99406, 99407	Reason For Visit
Tobacco use counseling: pregnant women	The USPSTF recommends that clinicians ask all pregnant women about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant women who use tobacco.	99406, 99407	Reason For Visit
Tobacco use interventions: children and adolescents	The USPSTF recommends that clinicians provide interventions, including education or brief counseling, to prevent initiation of tobacco use in school-aged children and adolescents.	99406, 99407	Reason For Visit
Tuberculosis screening: adults	The USPSTF recommends screening for latent tuberculosis infection in populations at increased risk.	86480*, 86481*, 86580*, 87116*	Z111, Z201

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

USPSTF A and B RECOMMENDATIONS

***In addition to modifier 33, a diagnosis to support the preventive service is required for services provided to Fee-For-Service beneficiaries.**

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	PREVENTIVE SERVICES	RECOMMENDED DIAGNOSIS
Vision screening: children	The USPSTF recommends vision screening at least once in all children ages 3 to 5 years to detect amblyopia or its risk factors.	Covered during preventive medicine services or a focused E/M visit.	Reason For Visit

Advisory Committee on Immunization Practices (ACIP) recommended vaccines

Follow current Medicaid policy for coverage of ACIP recommended vaccines including VFC.

Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine Recommended Preventive Health Care Services for Women (formerly known as Institute of Medicine)

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	HEALTHY MICHIGAN	PREVENTIVE CPT CODES
Screening for gestational diabetes (Recommendation 5.1)	Screening for gestational diabetes in pregnant women between 24 and 28 weeks of gestation and at the first prenatal visit for pregnant women identified to be at high risk for diabetes.	82950-82952	See USPSTF A and B RECOMMENDATIONS Above

Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine Recommended Preventive Health Care Services for Women (formerly known as Institute of Medicine)

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	HEALTHY MICHIGAN	PREVENTIVE CPT CODES
Human papillomavirus testing (Recommendation 5.2)	The addition of high-risk human papillomavirus DNA testing in addition to cytology testing in women with normal cytology results. Screening should begin at 30 years of age and should occur no more frequently than every 3 years.	87623-87625, G0476	See USPSTF A and B RECOMMENDATIONS Above
Counseling for sexually transmitted infections (Recommendation 5.3)	Annual counseling on sexually transmitted infections for sexually active women.	Covered during preventive medicine services or a focused E/M visit.	See USPSTF A and B RECOMMENDATIONS Above

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

<p>Counseling and screening for human immune-deficiency virus (Recommendation 5.4)</p>	<p>Counseling and screening for human immunodeficiency virus infection on an annual basis for sexually active women.</p>	<p>86689, 86701-86703, 87389-87391, 87534-87539, 87806, G0432-G0433, G0435</p> <p>Counseling covered during preventive medicine services or a focused E/M visit. Labs as indicated.</p>	<p>See USPSTF A and B RECOMMENDATIONS Above</p>
<p>Contraceptive methods and counseling (Recommendation 5.5)</p>	<p>The full range of Food and Drug Administration-approved contraceptive methods, sterilization procedures, and patient education and counseling for women with reproductive capacity.</p>	<p>11976, 11981, 11982, 11983, 57170, 58300, 58301, 58565, 58600, 58605, 58611, 58615, 58670, 58671, A4264, A4266, A4268, A4269, J1050, J7296, J7297, J7298, J7300, J7301, J7304, J7307, S4989</p> <p>Counseling covered during preventive medicine services or a focused E/M visit. Oral contraceptives are included as a pharmacy benefit.</p>	<p>The Medicaid program covers medically necessary contraceptive and counseling services for eligible beneficiaries.</p>
<p>Breastfeeding support, supplies and counseling (Recommendation 5.6)</p>	<p>Comprehensive lactation support and counseling and costs of renting breastfeeding equipment. A trained provider should provide counseling services to all pregnant women and to those in the postpartum period to ensure the successful initiation and duration of breastfeeding. (The ACA ensures that breastfeeding counseling is covered; however, the committee recognizes that interpretation of this varies.)</p>	<p>A4281-A4286, E0602-E0604 S9443</p>	<p>See USPSTF A and B RECOMMENDATIONS Above</p>

Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine

Recommended Preventive Health Care Services for Women (formerly known as Institute of Medicine)

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	HEALTHY MICHIGAN	PREVENTIVE CPT CODES
<p>Screening and counseling for interpersonal and domestic violence (Recommendation 5.7)</p>	<p>Screening and counseling for interpersonal and domestic violence. Screening and counseling involve elicitation of information from women and adolescents about current and past violence and abuse in a culturally sensitive and supportive manner to address current health concerns about safety and other current or future health problems.</p>	<p>Covered during preventive medicine services or a focused E/M visit.</p>	<p>See USPSTF A and B RECOMMENDATIONS Above</p>

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

Well-woman visit (Recommendation 5.8)	At least one well-women preventive care visit annually for adult women to obtain the recommended preventive services, including preconception and prenatal care. The committee also recognizes that several visits may be needed to obtain all necessary recommended preventive services, depending on a woman's health status, health needs, and other risk factors.	Covered during preventive medicine services.	Covered during preventive medicine services.
---------------------------------------	---	--	--

Early and Periodic Screening, Diagnosis and Treatment (EPSDT) services (19 and 20 year olds only)

Follow current Medicaid policy for coverage of EPSDT services.

Revision History

04/2014	<ul style="list-style-type: none"> - Added HCPCS codes: A4281-A4286 and E0602-E0604, to IOM's <i>Breastfeeding support, supplies and counseling preventive service</i>. - <i>Gestational diabetes mellitus screening</i> added to USPSTF January 2014
07/2014	<ul style="list-style-type: none"> - <i>Lung cancer screening</i> added to USPSTF December 2013 - <i>Hepatitis B screening: nonpregnant adolescents and adults</i> added to USPSTF May 2014 - <i>BRCA screening, counseling about</i> had an update to the USPSTF preventive service title. - <i>Dental caries prevention: preschool children</i> had an update to the USPSTF preventive service title. - Align <i>HIV screening</i> with current 2014 USPSTF version.
10/2014	<ul style="list-style-type: none"> - <i>Healthy diet counseling</i> had an update to the USPSTF preventive service title and description. - <i>Preeclampsia prevention: aspirin</i> added to USPSTF September 2014 - <i>Chlamydia screening: women</i> had an update to the USPSTF preventive service title and description. - <i>Gonorrhea screening: women</i> had an update to the USPSTF description. - <i>Sexually transmitted infections counseling</i> had an update to the USPSTF description.
02/2015	<ul style="list-style-type: none"> - Added 2015 HCPCS codes: 45346, 45388, 77063, 77085-77086, 87623-87625, 87806, G0464, G0472, G6019, G6022, G6024 - Deleted discontinued HCPCS codes: 45339, 45383, 87620-87622
10/2015	<ul style="list-style-type: none"> - <i>High blood pressure in adults: screening</i> added to USPSTF October 2015 - <i>Diabetes Screening</i> had an update to the USPSTF description.
01/2016	<ul style="list-style-type: none"> - Added HCPCS code: A4264 - Added 2016 HCPCS codes: 80081, 81528, G0297, J7297, J7298 - Deleted HCPCS code: G0464 as it is a Grade I. - Deleted discontinued HCPCS codes: G6019, G6022, G6024, J7302
04/2016	<ul style="list-style-type: none"> - <i>Depression screening: adults</i> had an update to the USPSTF preventive service description. - <i>Depression screening: adolescents</i> had an update to the USPSTF preventive service description. - Added HCPCS code: S9443 - Revised notes concerning Lactation support.
07/2016	<ul style="list-style-type: none"> - <i>Diabetes screening:</i> Added HCPCS codes: 82952 - <i>Hepatitis B screening: nonpregnant adolescents and adults:</i> Deleted HCPCS codes: 86707, 87341 - <i>Hepatitis B screening: pregnant women:</i> Deleted HCPCS codes: 80055, 80081, 86707, 87341 - <i>Osteoporosis screening: women:</i> Deleted HCPCS codes: 77085 and 77086. Added HCPCS codes: 76977, 77078, 77081 - <i>Syphilis screening: nonpregnant/pregnant persons</i> had an update to the USPSTF preventive service description.

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

	<ul style="list-style-type: none"> - Aspirin to prevent cardiovascular disease: men/women had an update to the USPSTF preventive service title and description. - Blood pressure screening in adults had an update to the USPSTF preventive service description. - Tobacco use counseling: nonpregnant/pregnant had an update to the USPSTF preventive service description. - Anemia screening: pregnant women removed from USPSTF A and B recommendation. - Iron supplementation in children: removed from USPSTF A and B recommendation. - Colorectal cancer screening had an update to the USPSTF preventive service description.
11/2016	<ul style="list-style-type: none"> - Tuberculosis screening: adults added to USPSTF September 2016 - Breastfeeding counseling had an update to the USPSTF preventive service title and description. - Statin preventive medication: adults ages 40–75 years with no history of CVD, 1 or more CVD risk factors, and a calculated 10-year CVD event risk of 10% or greater added to USPSTF November 2016
01/2017	<ul style="list-style-type: none"> - Deleted discontinued HCPCS codes: G0436, G0437, 77052, 77057 - Added 2017 HCPCS codes: G0499, 77067 - Added HCPCS code: G0476
04/2017	<ul style="list-style-type: none"> - Cholesterol abnormalities screening: men 35 and older removed from USPSTF A and B recommendation. - Cholesterol abnormalities screening: men younger than 35 removed from USPSTF A and B recommendation. - Cholesterol abnormalities screening: women 45 and older removed from USPSTF A and B recommendation. - Cholesterol abnormalities screening: women younger than 45 removed from USPSTF A and B recommendation. - High blood pressure in adults: screening removed from USPSTF A and B recommendation.
07/2017	<ul style="list-style-type: none"> - Deleted HCPCS code: 99402 and note - Preeclampsia: screening added to USPSTF April 2017 - Obesity screening: children and adolescents had an update to the USPSTF preventive service title and description. - The format of this document was updated to include information reflective of the coverage of preventive services for both Healthy Michigan Plan beneficiaries and Medicaid Fee-for-Service beneficiaries.
09/2017	<ul style="list-style-type: none"> - Vision acuity screening: children had an update to the USPSTF preventive service title and description.
01/2018	<ul style="list-style-type: none"> - Alcohol misuse: screening and counseling: Added 99408 and 99409 - Deleted discontinued HCPCS codes: G0202, Q9984 - Added 2018 HCPCS codes: 00812, J7296